



Southeast Tennessee HIV Community Planning Region, 2022



65 persons were newly diagnosed with HIV



25%
received a late diagnosis

→ Persons diagnosed with HIV Stage 3 (AIDS) within 12 months of a new HIV diagnosis are considered a **late diagnosis**, indicating delayed awareness of HIV status and missed opportunities for prevention efforts.

Most affected subpopulations by number of persons newly diagnosed with HIV


Non-Hispanic black men, MMS	13
Non-Hispanic white men, MMS	9
Non-Hispanic black men, HET	7
Non-Hispanic white men, MMS+IDU	6
Hispanic men, MMS	5
Non-Hispanic white women, IDU	4
Non-Hispanic white men, IDU	4

MMS: male-to-male sexual contact; HET: heterosexual contact; IDU: injection drug use

68% were linked to HIV care within 30 days of diagnosis

91% were linked within 365 days

1,726 persons were living with diagnosed HIV



62%
were virally suppressed

→ When HIV medication reduces the amount of HIV virus in a person's body to a low enough level (<200 c/mL), they are considered **virally suppressed** and will not transmit HIV through sexual contact.



For more HIV surveillance data visit: tn.gov/health/HIVData

Due to standard reporting delays, the data in this report represent all information reported to TDH through September 14, 2023. Source: Tennessee enhanced HIV/AIDS Reporting System (eHARS)